

(b) translating said nucleic acid sequence under conditions in which translation stalls at the 3' end of said nucleic acid sequence, forming a stalled translation complex comprising said protein; and

A12 (c) contacting said stalled translation complex with a puromycin-tag under conditions in which said puromycin-tag is covalently bonded to the C-terminus of said protein.

15. (Amended) The method of claim 14, wherein the tag of said puromycin-tag is attached to the 5'-hydroxy group of the puromycin.

---

A13 18. (Amended) The method of claim 14, wherein said translating step (b) is carried out in the substantial absence of at least one translation release factor.

---

A14 20. (Amended) The method of claim 14, wherein the tag of said puromycin-tag is a small molecule.

---

A15 22. (Amended) The method of claim 14, wherein the tag of said puromycin-tag is a detectable label.

---

A16 24. (Amended) The method of claim 14, wherein the tag of said puromycin-tag is a functional group.

25. (Amended) The method of claim 14, wherein said protein has a first functional group and the tag of said puromycin-tag is a second functional group and wherein said first functional group has a reactivity orthogonal to the reactivity of said second functional group.

A16

26. (Amended) The method of claim 14, wherein the tag of said puromycin-tag is a tether for attachment to a solid support.

---

28. (Amended) The method of claim 14, wherein the tag of said puromycin-tag is one member of a specific binding pair.

A17

29. (Amended) The method of claim 28, wherein said one member is a phenyl diboronic acid derivative.

30. (Amended) The method of claim 14, wherein a nucleotide sequence is positioned between the tag and the puromycin of said puromycin-tag.

---

In the Drawings:

Kindly amend Figure 3 as indicated in the accompanying marked up version of this figure, and replace original Figures 1-12 with the attached formal Figures 1-12.